arXiv:0704.2435v2 [nlin.SI] 1 May 2007

Integrable Nonautonomous Nonlinear Schrödinger Equations

Metin Gürses

Department of Mathematics, Faculty of Sciences

Bilkent University, 06533 Ankara - Turkey

February 1, 2008

Abstract

We show that a recently given nonautonomous nonlinear Schrodinger equation (NLSE) can be transformed into the autonomous NLSE.

1 Introduction

In [1] we have shown that integrable variable coefficient KdV type of equations, $q_t = f(x,t)\, q_3 + H(x,t,q,q_1)$ are transformable into autonomous integrable KdV type of equations, $q_t = q_3 + F(q,q_1)$. It seems that this assertion is valid in general. In a recent paper [2] an integrable nonautonomous NLSE is introduced

$$iQ_t + \frac{D(t)}{2}\, Q_{xx} + \sigma R(t)\, |Q|^2\, Q - [2\alpha(t)\, x + \frac{\Omega^2(t)}{2}\, x^2\,]\, Q = 0 \tag{1}$$

where the functions $R(t)$, $D(t)$, and $\Omega(t)$ representing the nonlinear, dispersion and the harmonic potential terms respectively are restricted to satisfy the following condition.

$$-\Omega^2(t)\, D(t) = \frac{d^2}{dt^2} \ln D(t) + R(t) \frac{d^2}{dt^2} \frac{1}{R(t)} - \frac{d}{dt} \ln D(t)\, \frac{d}{dt} \ln R(t) \tag{2}$$

They give a Lax pair of this equation where integrability implies both (1) and (2). The spectral parameter is a time dependent function and hence solitons obtained through the inverse scattering method have time dependent amplitudes and speeds. We shall first show that (1) is transformable to the standard NLSE and produce the soliton solutions given in [2] directly from the soliton solution of the NLSE.

Let $Q = \Lambda\, q(X,T)$ where

$$T = G(t), \quad X = F(x,t), \quad \Lambda = \Lambda(x,t) \tag{3}$$

Then Eq.(1) reduces to the standard NLSE

$$iq_T + q_{XX} + \sigma\, |q|^2\, q = 0, \tag{4}$$

where the following conditions must be satisfied.

$$i\Lambda_t + \frac{D(t)}{2}\,\Lambda_{xx} - 2\alpha(t)x\Lambda - \frac{\Omega^2(t)}{2}x^2\Lambda = 0, \tag{5}$$

$$i\Lambda F_t + \frac{D(t)}{2}[2\Lambda_x F_x + \Lambda F_{xx}] = 0, \tag{6}$$

$$\frac{D(t)}{2}\frac{F_x^2}{G_t} = 1, \quad \frac{R|\Lambda|^2}{G_t} = 1 \tag{7}$$

One can find a solution of the above equations. One solution is given as follows

$$\Lambda = r(t)e^{i\theta}, \quad \theta = -\frac{r_t}{D(t)r}\,x^2 + \alpha_1\,x + \alpha_2, \tag{8}$$

$$r^2 = 2r_0^4\frac{R(t)}{D(t)}, \tag{9}$$

where α_1 and α_2 are functions of t, r_0 is an arbitrary real constant and r is subjected to satisfy the equation

$$\frac{d}{dt}\left(\frac{1}{r(t)D(t)}\frac{dr}{dt}\right) - \frac{D(t)}{2}\left(\frac{2}{r(t)D(t)}\frac{dr}{dt}\right)^2 - \frac{\Omega^2(t)}{2} = 0 \tag{10}$$

When $r(t)$ given in (9) is used in (10) we obtain exactly the condition in (2). Furthermore we obtain $F = \frac{r^2}{r_0^2}\,x + F_1$ and

$$\alpha_1 = -2\frac{R(t)}{D(t)}\int\frac{\alpha(t)D(t)}{R(t)}dt, \tag{11}$$

$$\alpha_2 = -\frac{1}{2}\int D(t)\,{\alpha_1}^2 dt, \tag{12}$$

$$F_1 = -r_0^2\int \alpha_1(t)\,R(t)dt, \tag{13}$$

$$G(t) = 2r_0^4\int\frac{R^2(t)}{D(t)}\,dt \tag{14}$$

One soliton solution of the NLSE (4) was given by Zakharov and Shabat [3] which depends on the sign σ. For $\sigma = 1$ we have

$$q(X,T) = a \exp\left[i\left\{\frac{c}{2}(X-cT)+nT\right\}\right] \operatorname{sech}\left\{a(X-cT)/\sqrt{2}\right\} \tag{15}$$

where $4n = c^2 + 2a^2$. For $\sigma = -1$ we have

$$q(X,T) = \frac{1}{\sqrt{2}}\left[c - 2i\kappa \tanh \kappa(X-cT)\right]e^{-imt} \tag{16}$$

where $c^2 = 2m - 4\kappa^2$. Inserting the transformations (3) in the above Zakharov-Shabat solutions we obtain the solutions obtained by Serkin et al [2].

I would like to thank Atalay Karasu for discussions. This work is partially supported by the Turkish Academy of Sciences.

References

[1] M. Gürses and A. Karasu, J. Math. Phys. **36**, 3485 (1995).

[2] V.N. Serkin, A. Hasegawa, and T.L. Belyaeva, Phys. Rev. Lett. **98**, 074102 (2007).

[3] V.E. Zakharov and A.B. Shabat, Sov. Phys. JETP., **37**, 823 (1973).